



TITLE:

Diversity and evolution of pollinator rewards and protection by *Macaranga* (Euphorbiaceae) bracteoles

AUTHOR(S):

Yamasaki, Eri; Kawakita, Atsushi; Sakai, Shoko

CITATION:

Yamasaki, Eri ...[et al]. Diversity and evolution of pollinator rewards and protection by *Macaranga* (Euphorbiaceae) bracteoles. *Evolutionary Ecology* 2015, 29(3): 379-390

ISSUE DATE:

2015-05-16

URL:

<http://hdl.handle.net/2433/200785>

RIGHT:

The final publication is available at Springer via <http://dx.doi.org/10.1007/s10682-014-9750-7>; The full-text file will be made open to the public on 16 May 2016 in accordance with publisher's 'Terms and Conditions for Self-Archiving'; This is not the published version. Please cite only the published version.; この論文は出版社版ではありません。引用の際には出版社版をご確認ください。

1
2 Diversity and evolution of pollinator rewards and protection by *Macaranga* (Euphorbiaceae)
3 bracteoles
4
5 Eri Yamasaki
6 Center for Ecological Research, Kyoto University
7 Hirano 2-509-3, Otsu, Shiga 520-2113, Japan
8 Institute of Evolutionary Biology and Environmental Studies, University of Zurich
9 Winterthurerstrasse 190, CH-8057 Zurich, Switzerland
10 yamasakiebi@gmail.com
11 Atsushi Kawakita
12 Center for Ecological Research, Kyoto University
13 Hirano 2-509-3, Otsu, Shiga 520-2113, Japan
14 Shoko Sakai
15 Center for Ecological Research, Kyoto University
16 Hirano 2-509-3, Otsu, Shiga 520-2113, Japan
17
18
19 Running title:
20 Diversity and evolution of bracteoles in *Macaranga* (Euphorbiaceae)
21
22 Key words:
23 Ant-plants, Euphorbiaceae, Extrafloral nectary, Hemipteran pollination, *Macaranga*, Thrips
24 pollination
25

- 26 Total word count: 3371
- 27 Introduction: 997
- 28 Materials and Methods: 847
- 29 Results: 472
- 30 Discussion: 708
- 31 Number of cited references: 48
- 32
- 33 Number of figures: 4
- 34 Number of tables: 1
- 35
- 36 Items to be published as online appendix:
- 37 Supplemental Table 1. Inflorescence/floral traits assessed on herbarium specimens.
- 38 Supplemental Figure 1. A female inflorescence of *Macaranga vedeliana*.

Abstract

Flowering plants have modified their floral organs in remarkably diverse ways to optimize their interaction with pollinators. Although floral organs represent a major source of floral diversity, many plants also use extrafloral organs, such as bracts and bracteoles, in interacting with pollinators; however, the evolutionary dynamics of non-floral organs involved in pollination are poorly studied. The genus *Macaranga* is characterized by protective mutualisms with ants that potentially interfere with pollinators on flowers. *Macaranga* flowers lack perianths and, notably, bracteoles serve the dual function of rewarding pollinators and protecting them from guarding ants; in one group of species, bracteoles provide a nectar reward to generalist pollinators, while in another group, bracteole “chambers” protect thrips or hemipteran pollinators that use these structures as feeding and breeding sites. We examined the diversity and evolutionary dynamics of inflorescence morphology in *Macaranga*, focusing on bracteoles. We recognized three inflorescence types based on examination of herbarium materials: Discoid-gland, which possess disc-shaped glands on the bracteole surfaces (including all the generalist-pollinated species); Enclosing, in which bracteoles cover flowers (including all the thrips- and hemipteran-pollinated species); and Inconspicuous, in which bracteoles are small, narrow or absent. Ancestral state reconstruction indicated that inflorescence morphologies have changed multiple times in the genus. These findings suggest that morphological changes in non-floral characters (bracteoles) of *Macaranga* species have occurred as frequently as in the floral structures of many flowering plants. The multiple evolutions of the Enclosing bracteoles, which protect pollinators, might have been facilitated by pollination interference from mutualistic ants.

62 **Introduction**

63 Flowering plants exhibit enormously diverse floral traits, many of which are useful for animal
64 pollination (Ollerton et al. 2011). For example, showy perianths and odors can efficiently
65 attract pollinators, and nectars, oils, resins, and other floral secretions can reward pollinators
66 (Fægri and van der Pijl 1979). Therefore, biologists have studied how flower visitors
67 influence the evolution of floral traits in various plant groups by combining phylogenetic
68 relationships among focal plants, and their floral traits and pollination systems (Johnson et al.
69 1998; Beardsley et al. 2003; Whittall and Hodges 2007; Wilson et al. 2007; Okuyama et al.
70 2008; Kawakita and Kato 2009; Sakai et al. 2013). Overall, evidence suggests that the same
71 pollination systems have evolved repeatedly among plant groups, altering floral traits and
72 confirming their flexibility.

73 Although both floral and extrafloral organs are used for attracting and/or rewarding
74 pollinators, floral organs such as petals, sepals, and nectaries within flowers play more
75 important roles in pollination than extrafloral organs in most flowering plants. However, in
76 some plant species, especially in those which have lost conspicuous petals or sepals,
77 extrafloral organs are more important for interactions with pollinators; in most of these cases,
78 bracts and bracteoles play major roles (Ehrenfeld 1979; Armbruster 1997; Wragg and
79 Johnson 2011; Bröderbauer et al. 2012). The roles of bracts or bracteoles have been well
80 described in the pollination systems of the *Araceae*, and in the *Euphorbia*, *Dalechampia*
81 (*Euphorbiaceae*), and some *Cyperus* (*Cyperaceae*) species (Ehrenfeld 1979; Armbruster
82 1997; Wragg and Johnson 2011; Bröderbauer et al. 2012), but it is unclear whether bracts or
83 bracteoles in these plants change as flexibly as floral organs, or are under similar
84 evolutionary constraints. Some studies have suggested frequent changes in pollination
85 systems without large morphological changes; Armbruster (1993) and Armbruster (1997)
86 reported frequent pollinator reward shifts between resin and pheromone precursors or pollen

in *Dalechampia*, while the structure of bracts, the primary organ used to attract pollinators, remained consistent. In Araceae, evolution of pollinator-trapping chambers that enclose the spadices has occurred at least 10 times from inflorescences in which bracts do not cover the spadices completely. However, bract shape does not change drastically in many species in which bracts surround the spadices and form a chamber around the flowers (Bröderbauer et al. 2012).

It has been said that pollinators play important roles in driving plant traits involved in pollination (Whittall and Hodges 2007; van der Niet and Johnson 2012). However, flowers are visited not only by pollinators but also by unfavorable animals, such as florivores, worthless pollinators, or pollinator-excluding animals (primarily ants) (Willmer and Stone 1997; Irwin et al. 2004). These unfavorable visitors can also have a selective effect on floral/inflorescence traits; for example, long corollas exclude insects with short mouthparts, and pendulous flowers exclude all but hovering insects and those with strong legs (Proctor et al. 2006).

In this study, we investigated the evolution of inflorescence morphologies in the dioecious tree genus *Macaranga* (Euphorbiaceae), distributed in tropical to subtropical regions (Whitmore 2008). Recent studies have reported two distinct bracteole morphologies in the genus associated with different pollination systems. The first consists of paddle-shaped bracteoles with one to several disc-shaped nectaries on the adaxial (upper) surfaces (Fig. 1a; Yamasaki et al. 2013a). Pollinator insects visit the inflorescences in search of nectar secreted from the nectaries. The number and size of the bracteoles do not differ between male and female inflorescences. The second consists of bracteoles enclosing flowers associated with pollination by thrips, *Dolichothrips* spp. (Phlaeothripidae) or by hemipterans of the Anthocoridae and Miridae (Fig. 1b; Moog et al. 2002; Ishida et al. 2009; Fiala et al. 2011). The pollinators breed in bracteole chambers and feed on trichome- and/or ball-shaped

nectaries on the adaxial surfaces of the bracteoles. Although the shape is similar between male and female inflorescences, the bracteoles are stouter and fewer in number in female than in male inflorescences (Davies and Ashton 1999).

Macaranga is well known for its protective mutualisms with ants. Most of the ca. 260 species in the genus possess extrafloral nectaries and/or food bodies (nourishing small particles) on leaves and potentially attract ants that can exclude herbivores from flowers (Whalen and Mackay 1988; Fiala and Maschwitz 1991; Mackay and Whalen 1991; Whitmore 2008). On the other hand, ca. 30 species offer not only foods for ants but also nesting sites for them (Davies et al. 2001). In this paper, we call these species “ant-plants.” The ant-plant *Macaranga* species are intensely protected by the almost species-specific resident ants, compared to species that are facultatively protected by ants attracted to extrafloral nectaries and/or food bodies (Quek et al. 2004; Itioka 2005). While ant-plant *Macaranga* species are pollinated by *Dolichothrips* spp. (Fiala et al. 2011), our previous study indicated that pollinator thrips are not excluded by the guard ants (Yamasaki et al. 2013b). This may be because pollinators are protected from the guard ants by bracteoles (Fiala et al. 2011; Yamasaki et al. 2013b); although ant guards are often seen on the surface of the inflorescences, we have never observed the guard ants crawling into the chambers, where pollinator thrips spend most of their lifetime, maybe because the gaps of the bracteoles are too narrow for the ants.

While some previous studies have examined the evolution of inflorescence morphology in *Macaranga* (Davies 2001; Fiala et al. 2011), they mainly focused on the sister sections *Pachystemon* and *Pruinosae*, which have flower-enclosing bracteoles, no clear overview of the evolutionary pattern in the genus has been elucidated. The aim of this study was to examine whether inflorescence morphologies have changed repeatedly and, if so, whether the repeated evolution is related to interacting animals such as pollinators and

bodyguard ants in *Macaranga*. For this purpose, we first examined interspecific variation in inflorescence morphologies and estimated pollination systems of each morphology type by examining the inflorescence characteristics and previous pollination studies in *Macaranga*. Next, we mapped the inflorescence types on a molecular phylogenetic tree and estimated ancestral inflorescence morphologies to examine whether inflorescence morphologies have repeatedly changed in *Macaranga*.

Materials and Methods

Observation of inflorescence/floral morphologies

We observed the inflorescences of dry specimens of 53 taxa in the genus *Macaranga* (52 species and one variety) in herbaria (Royal Botanic Gardens, Kew (K), Leiden Naturalis Biodiversity Center (L), Kyoto University (KYO) and Forest Department, Sarawak (SAR)). We also obtained morphological data for *Macaranga lamellata* and *Macaranga umbrosa* from Fiala et al. (2011). We recorded (1) the presence/absence of disc-shaped glands on bracteole surfaces, (2) internode distances between adjacent bracteoles, (3) length, and (4) width of bracteoles in male specimens, and (5) style length in female specimens (Fig. 2). Traits (1)–(4) were not measured in female specimens because those with bracteoles were absent in many of the species, while most of the bracteoles in male inflorescences remained. For each trait, we looked at two to five samples from each of one to five specimens. For trait (1), we judged disc-shaped glands to be present when at least one bracteole possessed them, and we determined that specific taxa possessed the glands if they occurred in at least one specimen. The presence/absence of the glands was consistent among specimens in most species, with the exception of *Macaranga denticulata*. Some specimens of this species lack disc-shaped glands on their bracteoles, but in all other respects, their bracteoles were similar to those of other specimens. Thus, we regarded it as a species with disc-shaped glands. For

quantitative traits (2)–(5), average values were calculated for each specimen and averaged across specimens to obtain species values.

To determine which traits to use in categorizing inflorescence morphologies, we examined the differences in inflorescence morphologies among the species by principal component analysis (PCA) using *Z*-score standardized values of the four quantitative traits (traits (2)–(5)). Thirty-two taxa in which all four variables were available were included in the analysis. We used the *prcomp* function in R 3.0.2 (R Development Core Team 2013). The first principal component (PC 1) clearly separated species that did not contain disc-shaped glands into two groups, and bracteole length and width were major components (see Results). Therefore, we classified all 55 taxa into three inflorescence types based on the presence/absence of disc-shaped glands, and bracteole shape and size (see Results). Style length was not used because we were unable to measure it in many species, mostly due to a lack of specimens containing flowering female inflorescences.

Molecular phylogenetic analysis

We constructed a molecular phylogeny based on the DNA sequence and indel data on one plastid (*trnL-F*) and three nuclear markers (ITS, *ncpGS*, and *phyC*) of 59 taxa in the genus *Macaranga* using species of the related genus *Mallotus* (Euphorbiaceae) as outgroups (*Mallotus griffithianus*, *Mallotus claoxyloides*, and *Mallotus paniculatus*). Alignment and indel data for all taxa other than *Macaranga sinensis* were acquired from Kulju et al. (2007); those of *M. sinensis* were obtained via the following procedures. First, DNA was extracted from silica gel-dried leaves following a modified CTAB procedure (Doyle and Doyle 1987; Okuyama and Kawakita 2012). Regions were amplified by different primer pairs (Kulju et al. 2007). Polymerase chain reaction (PCR) amplifications were carried as follows: initial denaturation step at 94°C for 5 min; 30 cycles of 94°C for 30 s, 50°C for 30 s, and 72°C for 1

min; and a final extension at 72°C for 7 min. Products were sequenced on an ABI 3100 automated sequencer using BigDye chain termination chemistry (Applied Biosystems, Foster City, CA, USA), and obvious sequence errors were manually corrected using MEGA 5.05. Indel information was incorporated based on Kulju et al. (2007). A phylogenetic tree was constructed by Bayesian inference methods using MrBayes version 3.2 (Ronquist and Huelsenbeck 2003). The substitution models were chosen separately for each marker based on Akaike Information Criterion using Kakusan 4 (Tanabe 2011) as follows: GTR+G for *trnL* and ITS and HKY85+G for *ncpGS* and *phyC*. We employed the binary model for gap characters (Ronquist and Huelsenbeck 2003). Two independent runs for four Markov chains were conducted for four million generations, and the tree was sampled every 100 generations. The first 4001 trees were discarded as burn-in. A majority-rule consensus tree was constructed from the remaining 36000 trees.

Reconstruction of ancestral inflorescence morphologies

To investigate how inflorescence type has shifted in the genus, we estimated ancestral inflorescence morphologies (the inflorescence morphology of each node) using 1000 trees randomly chosen from the Bayesian phylogenetic analysis. Inflorescence morphologies of the outgroups are not included in the analysis. The Markov chain Monte Carlo method (Ronquist 2004) was applied for this analysis using the BayesMultiState function in BayesTraits (Pagel and Meade 2006). The reverse jump hyper prior function with a gamma prior of 0, 10, 0, 10 (minimum and maximum of priors for both mean and variance parameters) was used. Rate deviation was set to 50 to achieve the recommended 20–40% acceptance rates. Inflorescence morphology types were treated as categorical variables with three states (Fig. 1; see Results). Alternative ancestral states were compared on 49 nodes with >0.5 posterior probabilities using the ‘fossil’ (fixing) command. Because harmonic means in the analyses can be unstable

(Pagel and Meade 2006), we ran the analyses five times to check the consistency of the results. Bayes Factors ($2 [\log(\text{harmonic mean of model 1}) - \log(\text{harmonic mean of model 2})]$) >5 indicated strong evidence of support for the best model (Pagel et al. 2004).

Results

Observation of inflorescence/floral morphologies

Based on the PCA results, inflorescence types were classified into three categories using bracteole shape and size and the presence/absence of disc-shaped glands: Discoid-gland, Inconspicuous, and Enclosing (Fig. 1; Supplemental Table 1). In the PCA results, the first and second principal components (PC 1 and PC 2) contributed 54.1% and 24.6% of the total variance of the measured data, respectively (Table 1). Bracteole size (length and width) and style length had substantial loading in PC 1 (Table 1, Fig. 3). PC 1 distinctly separated species not possessing disc-shaped glands into two groups: one with relatively large bracteoles and short styles, and one with small bracteoles and long styles. PC 2 mainly represented internode distances between bracteoles and bracteole length, and some species with disc-shaped glands had extremely low values. The two groups distinguished by PC 1 were not represented by PC 2.

The three categories are defined as follows based on the PCA and presence/absence of disc-shaped glands:

(1a) Disc-shaped glands on the bracteole surfaces presentDiscoid-gland

(1b) Disc-shaped glands on the bracteole surfaces absent (2)

(2a) Bracteoles very small (length and width < 1.6 mm) or narrow (length/width > 1.8), or absent.....Inconspicuous

(2b) Bracteoles relatively large, enclosing flower clusters.....Enclosing

We also observed fresh inflorescences of 11 species in which all three inflorescence types

were represented. The shape did not differ between dry specimens and fresh inflorescences, although the size was smaller in the dry specimens. All species mainly visited by thrips or hemipterans were of the Enclosing type (Moog et al. 2002; Ishida et al. 2009; Fiala et al. 2011; Figs. 3 and 4), as were the ant-plant species (Davies 2001; Davies et al. 2001; Fig. 4). *M. sinensis*, pollinated by generalist insects attracted to the disc-shaped nectaries on bracteoles (Yamasaki et al. 2013a), and *M. denticulata* and *Macaranga indica*, whose male inflorescences are mainly visited by generalist insects (bees, flies, wasps, and beetles) (Fiala et al. 2011), were of the Discoid-gland type (Fig. 4).

Molecular phylogenetic analysis

Molecular analysis revealed two well-supported basal clades (B1 and B2) and three crown clades (C1, C2, and C3), as in Kulju et al. (2007), who analyzed the phylogeny by Bayesian and maximum parsimony methods (Fig. 4). While the Bayesian tree in Kulju et al. (2007) united the C1 and C2 clades, our tree united the C2 and C3 clades, as in the maximum parsimony tree in Kulju et al. (2007).

Reconstruction of ancestral bracteole morphologies

All observed species in basal clades B1 and B2 were classified into the Inconspicuous category (Fig. 4). Conversely, we detected all three inflorescence types in the crown clades. No inflorescence type was determined to be monophyletic. Among the 49 focal nodes, six, five, and 14 nodes were strongly supported as Inconspicuous, Discoid-gland, and Enclosing types, respectively (Bayes Factors > 5; Fig. 4). When Bayes factor >5 is considered to indicate a significant occurrence of the focal inflorescence type, ancestral state reconstruction indicated at least two shifts from the Discoid-gland to the Enclosing type and at least one shift from the Enclosing to the Inconspicuous type (Fig. 4, Table 2).

262

263 Discussion

264 All *Macaranga* species in this study were classified into three inflorescence types based on
265 bracteole morphological characteristics: Discoid-gland, Inconspicuous, and Enclosing (Fig. 1,
266 see Results for detailed characteristics). Considering the morphology and existing pollination
267 studies in *Macaranga*, we propose that the three types of inflorescence morphology are
268 related to different pollination systems. Because this study lacks empirical pollination surveys
269 for many of the species, we cannot yet test this hypothesis. However, it is often said that
270 floral morphologies are generated by functional groups of similar pollinators (Fægri and van
271 der Pijl 1979; Fenster et al. 2004). Wind may contribute, at least in part, to pollination in the
272 Inconspicuous species, which have exposed flowers, inflorescences that penetrate through the
273 leaf mass, and sometimes extremely long (up to 5 cm) styles (Bullock 1994; Culley et al.
274 2002). We observed that *Macaranga vedeliana*, with Inconspicuous inflorescences (not
275 included in this study), did not secrete nectar and were wind-pollinated (Yamasaki et al.
276 unpublished data; Supplemental Figure 2).

277 In Discoid-gland species, the glands on bracteoles, which are located adjacent to flowers,
278 may attract insects for pollination in the manner of *M. sinensis*, which is pollinated by
279 generalist insects that forage on the disc-shaped nectaries (Yamasaki et al. 2013a). Similarly,
280 male inflorescences of two other Discoid-gland species, *M. denticulata* and *M. indica*, are
281 visited by generalist insects such as bees, wasps, flies, and beetles (Fiala et al. 2011). On the
282 other hand, the nectaries on bracteoles may also attract ants, which may exclude pollinators
283 or contribute pollination. Further experimental studies are needed to confirm whether the
284 nectaries on the bracteoles attract pollinators and how the plants avoid disadvantage from
285 ants if the glands also attract ants.

286 All thrips- or hemipteran-pollinated species (Moog et al. 2002; Ishida et al. 2009; Fiala et

al. 2011) were classified as Enclosing species (Fig. 4). Pollination by wind or generalist insects such as bees and flies is unlikely for this inflorescence type because such pollen vectors are impeded by the bracteoles. In addition to providing chambers for pollinators, the bracteoles of the Enclosing type may also physically protect pollinators against their natural enemies, such as ants (Fiala et al. 2011; Yamasaki et al. 2013b).

Ancestral state reconstruction indicated at least two shifts from the Discoid-gland to the Enclosing type and at least one shift from the Enclosing to the Inconspicuous type (Fig. 4, Table 2). Additionally, while direction is ambiguous, among the three inflorescence types, other shifts seem to have occurred several times (Fig. 4, Table 2). The present study is the first to describe multiple drastic evolutionary changes in extrafloral plant organs.

Repeated evolution of the Enclosing inflorescence type might be due to pollination interference by ants. Because bracteoles can physically separate ants and pollinators on the inflorescences, pollination interference by ants may have selected for the evolution of flower-covering bracteoles that eliminate ants from flowers. Therefore, the flower-enclosing bracteoles in the Enclosing inflorescences may act as “barriers” against pollination-interfering animals (Santamaría and Rodríguez-Gironés 2007). To explore these possibilities, the functions of bracteoles should be examined in Enclosing species from various clades. It would also be instructive to investigate why species with other inflorescence types do not need to exclude guard ants from their flowers. Because ant-plants are likely to have occurred more than once in an Enclosing-type clade (Blattner et al. 2001; Davies et al. 2001), the evolution of ant-plants and the development of the Enclosing inflorescences may be related. Not only mutualisms with ants but also the lack of perianths may be related to the multiple evolutions of the Enclosing type; because *Macaranga* lacks perianths that visually attract generalist pollinators such as bees, specialization for pollination by thrips or hemipterans, which infest the flowers of various plants (Lewis 1973; Wheeler

312 2001), may have occurred relatively easily.

313 In conclusion, our study found high variability in bracteole morphology compared with
314 that of flowers. *Macaranga* species have repeatedly evolved unusual traits involved in
315 pollination, including flower-enclosing bracteoles. The repeated evolution of
316 flower-enclosing bracteoles may indicate a recurring need for floral barriers against
317 bodyguard ants. The genus *Macaranga* may be a good model for studying the factors
318 underlying the acquisition of floral barriers.

319

320 *Acknowledgements*

321 We are grateful to herbaria of Royal Botanic Gardens, Kew (K), Leiden Naturalis
322 Biodiversity Center (L), Kyoto University (KYO) and Department of Forestry, Sarawak
323 (SAR) for providing *Macaranga* specimens. This study was financially supported by the
324 Research Institute for Humanity and Nature (Project D-04), Grants-in-Aid No. 21570028
325 from the Ministry of Education, Culture, Sports, Science and Technology, Japan, to S.S.,
326 Grants-in-Aid for JSPS Fellows from the Japan Society for the Promotion of Science (No.
327 24-1464 to E.Y.), and Kyoto University Global COE program “Formation of a strategic base
328 for biodiversity and evolutionary research: from genome to ecosystem”.

329 **References**

- 330 Armbruster WS (1993) Evolution of plant pollination systems: hypotheses and tests with the
331 Neotropical vine *Dalechampia*. *Evolution* 47:1480–1505
332
- 333 Armbruster WS (1997) Exaptations link evolution of plant-herbivore and plant-pollinator
334 interactions: a phylogenetic inquiry. *Ecology* 78:1661–1672
335
- 336 Beardsley PM, Yen A, Olmstead RG (2003) AFLP phylogeny of *Mimulus* section
337 *Erythranthe* and the evolution of hummingbird pollination. *Evolution* 57:1397–1410
338
- 339 Blattner FR, Weising K, Bänfer G, et al. (2001) Molecular analysis of phylogenetic
340 relationships among myrmecophytic *Macaranga* species (Euphorbiaceae). *Mol Phylogenet*
341 *Evol* 19:331–44
342
- 343 Bröderbauer D, Diaz A, Weber A (2012) Reconstructing the origin and elaboration of
344 insect-trapping inflorescences in the Araceae. *Am J Bot* 99:1666–1679
345
- 346 Bullock SH (1994) Wind pollination of Neotropical dioecious trees. *Biotropica* 26:172–179
347
- 348 Culley TM, Weller SG, Sakai AK (2002) The evolution of wind pollination in angiosperms.
349 *Trends Ecol Evol* 17:361–369
350
- 351 Davies SJ (2001) Systematics of *Macaranga* Sects. *Pachystemon* and *Pruinosae*
352 (Euphorbiaceae). *Harvard Pap Bot* 6:371–448
353

- 354 Davies SJ, Ashton PS (1999) Phenology and fecundity in 11 sympatric pioneer species of
355 *Macaranga* (Euphorbiaceae) in Borneo. *Am J Bot* 86:1786–1795
356
- 357 Davies SJ, Lum SKY, Chan R, Wang LK (2001) Evolution of myrmecophytism in western
358 Malesian *Macaranga* (Euphorbiaceae). *Evolution* 55:1542–1559
359
- 360 Doyle JJ, Doyle JL (1987) A rapid DNA isolation procedure for small quantities of fresh leaf
361 tissue. *Phytochem Bull* 19:11–15
362
- 363 Ehrenfeld JG (1979) Pollination of three species of *Euphorbia* subgenus *Chamaesyce*, with
364 special reference to bees. *Am. Midl. Nat.* 101:87–98
365
- 366 Fægri K, van der Pijl L (1979) The principles of pollination ecology. Pergamon Press, Oxford
367
- 368 Fenster CB, Armbruster WS, Wilson P, Dudash MR, Thomson JD (2004) Pollination
369 syndromes and floral specialization. *Annu Rev Ecol Evol Syst* 35:375–403
370
- 371 Fiala B, Maschwitz U (1991) Extrafloral nectaries in the genus *Macaranga* (Euphorbiaceae)
372 in Malaysia: comparative studies of their possible significance as predispositions for
373 myrmecophytism. *Biol J Linn Soc* 44:287–305
374
- 375 Fiala B, Meyer U, Hashim R, Maschwitz U (2011) Pollination systems in pioneer trees of the
376 genus *Macaranga* (Euphorbiaceae) in Malaysian rainforests. *Biol J Linn Soc* 103:935–953
377
- 378 Irwin RE, Adler LS, Brody AK (2004) The dual role of floral traits: pollinator attraction and

- 379 plant defense. Ecology 85:1503–1511
- 380
- 381 Ishida C, Kono M, Sakai S (2009) A new pollination system: brood-site pollination by flower
- 382 bugs in *Macaranga* (Euphorbiaceae). Ann Bot 103:39–44
- 383
- 384 Itioka T (2005) Diversity of anti-herbivore defences in *Macaranga*. In: Roubik DW, Sakai S,
- 385 Karim AAH (eds) Pollination ecology and the rain forest: Sarawak studies. Springer, New
- 386 York, pp 158–171
- 387
- 388 Johnson S, Linder H, Steiner K (1998) Phylogeny and radiation of pollination systems in
- 389 *Disa* (Orchidaceae). Am J Bot 85:402–411
- 390
- 391 Kawakita A, Kato M (2009) Repeated independent evolution of obligate pollination
- 392 mutualism in the *Phyllanthaceae–Epicephala* association. Proc Biol Sci 276:417–426
- 393
- 394 Kulju KKM, Sierra SEC, Draisma SGA, Samuel R, van Welzen PC (2007) Molecular
- 395 phylogeny of *Macaranga*, *Mallotus*, and related genera (Euphorbiaceae s.s.): insight from
- 396 plastid and nuclear DNA sequence data. Am J Bot 94(10):1726–1743
- 397
- 398 Lewis T (1973) Thrips: their biology, ecology and economic importance. Academic Press,
- 399 London
- 400
- 401 Mackay DA, Whalen MA (1991) Some associations between ants and euphorbs in tropical
- 402 Australia. In: Huxley C, Cutler DF (eds) Ant–plant interactions. Oxford Univ. Press, Oxford,
- 403 pp 238–249

404

405 Moog U, Fiala B, Federle W, Maschwitz U (2002) Thrips pollination of the dioecious ant
406 plant *Macaranga hullettii* (Euphorbiaceae) in Southeast Asia. *Am J Bot* 89(1):50–59

407

408 Okuyama U, Kawakita A (2013) DNA extraction protocol from plants (altered CTAB
409 method). In: Kawakita A, Okuyama U, ed. *Biology of Species Interactions*. Bun-ichi Sogo
410 Shuppan Co., Tokyo pp 273–278 [In Japanese]

411

412 Okuyama Y, Pellmyr O, Kato M (2008) Parallel floral adaptations to pollination by fungus
413 gnats within the genus *Mitella* (Saxifragaceae). *Mol Phylogenet Evol* 46:560–575

414

415 Ollerton J, Winfree R, Tarrant S (2011) How many flowering plants are pollinated by
416 animals? *Oikos* 120:321–326

417

418 Pagel M, Meade A (2006) Bayesian analysis of correlated evolution of discrete characters by
419 reversible-jump markov chain monte carlo. *Am Nat* 167:808–825

420

421 Pagel M, Meade A, Barker D. (2004) Bayesian estimation of ancestral character states on
422 phylogenies. *Syst Biol* 53:673–684

423

424 Proctor M, Yeo P, Lack A (2006) *The natural history of pollination*. Harper Collins, London

425

426 Quek S-P, Davies SJ, Itino T, Pierce NE (2004) Codiversification in an ant-plant mutualism:
427 stem texture and the evolution of host use in *Crematogaster* (Formicidae: Myrmicinae)
428 inhabitants of *Macaranga* (Euphorbiaceae). *Evolution* 58:554–570

429

430 R Development Core Team (2013) R: A language and environment for statistical computing.

431 R Foundation for Statistical Computing, Vienna, Austria. URL <http://www.R-project.org/>

432

433 Ronquist F (2004) Bayesian inference of character evolution. *Trends Ecol Evol* 19:475–481

434

435 Ronquist F, Huelsenbeck JP (2003) MrBayes 3: Bayesian phylogenetic inference under
436 mixed models. *Bioinformatics* 19:1572–1574

437

438 Sakai S, Kawakita A, Ooi K, Inoue T (2013) Variation in the strength of association among
439 pollination systems and floral traits: Evolutionary changes in the floral traits of Bornean
440 gingers (Zingiberaceae). *Am J Bot* 100:546–555

441

442 Santamaría L, Rodríguez-Gironés MA (2007) Linkage rules for plant–pollinator networks:
443 trait complementarity or exploitation barriers? *PLoS Biol* 5:e31

444

445 Tanabe AS (2011). Kakusan4 and Aminosan: two programs for comparing nonpartitioned,
446 proportional and separate models for combined molecular phylogenetic analyses of
447 multilocus sequence data. *Mol Ecol Resources* 11:914–921

448

449 Van der Niet T, Johnson SD (2012) Phylogenetic evidence for pollinator-driven
450 diversification of angiosperms. *Trends Ecol Evol* 27:353–361

451

452 Whalen MA, Mackay DA (1988) Patterns of ant and herbivore activity on five understory
453 euphorbiaceous saplings in submontane Papua New Guinea. *Biotropica* 20:294–300

454

455 Wheeler AG (2001) Biology of the plant bugs (Hemiptera: Miridae): pests, predators,
456 opportunists. Cornell University Press, New York

457

458 Whittall JB, Hodges SA (2007) Pollinator shifts drive increasingly long nectar spurs in
459 columbine flowers. Nature 447:706–710

460

461 Whitmore TC (2008) The genus *Macaranga*—a prodromus. Kew Publishing, Kew

462 Whittall JB, Hodges S (2007) Pollinator shifts drive increasingly long nectar spurs in
463 columbine flowers. Nature 447:706–710

464

465 Willmer PG, Stone GN (1997) How aggressive ant-guards assist seed-set in Acacia flowers.
466 Nature 388:165–167

467

468 Wilson P, Wolfe AD, Armbruster WS, Thomson JD (2007) Constrained lability in floral
469 evolution: counting convergent origins of hummingbird pollination in *Penstemon* and
470 *Keckiella*. New Phytol 176:883–890

471

472 Wragg PD, Johnson SD (2011) Transition from wind pollination to insect pollination in
473 sedges: experimental evidence and functional traits. New Phytol 191:1128–1140

474

475 Yamasaki E, Inui Y, Sakai S (2013b) Production of food bodies on the reproductive organs of
476 myrmecophytic *Macaranga* species (Euphorbiaceae): effects on interactions with herbivores
477 and pollinators. Plant Species Biol (in press)

478

- 479 Yamasaki E, Kawakita A, Sakai S (2013a) Modified leaves with disc-shaped nectaries of
480 *Macaranga sinensis* (Euphorbiaceae) provide reward for pollinators. Am J Bot 100:628–32
481
482

Figure Legends

Fig. 1. Examples of species exhibiting the three inflorescence types, categorized based on bracteole morphology: (a) *Macaranga sinensis* (Discoid-gland type), (b) *Macaranga gigantea* (Enclosing type), and (c) *Macaranga coriacea* (Inconspicuous type) (not included in the phylogeny). Scale bar = 1 cm.

Fig. 2. The inflorescence/floral traits investigated. See (1)–(5) in the *Observation of inflorescence/floral morphologies* section of Materials and Methods.

Fig. 3. Scatterplot of the first and second principal components (PC 1 and PC 2) of a principal components analysis (PCA) using four inflorescence and floral traits (internode distances between adjacent bracteoles, lengths and widths of bracteoles, internodes between two bracteoles, and style lengths). Different colors indicate inflorescence types (see text for classifications). Species visited mainly by thrips or hemipterans and those visited by other insects are indicated by diamond and triangle symbols, respectively. The information on flower visitors was obtained from Moog et al. (2002), Ishida et al. (2009), Fiala et al. (2011), and Yamasaki et al. (2013).

Fig. 4. Phylogenetic tree of *Macaranga* species constructed with the Bayesian inference method. The pies indicate estimated ancestral morphologies by Bayesian inference methods. Bayes factors >5 are indicated with an asterisk. Numbers above branches indicate posterior probabilities of the branches obtained from reconstruction of the tree (>0.5). The mapped information, with the exception of inflorescence morphologies, was obtained from the following literature: main flower visitors, Moog et al. (2002), Ishida et al. (2009), Fiala et al.

508 (2011), and Yamasaki et al. (2013a); whether the species are ant-plants, Davies (2001) and
509 Davies et al. (2001). Clade grouping (B1, B2, C1, C2, and C3) was done according to Kulju
510 *et al.* (2007).
511



512 **Table 1.** The proportion of variance and factor loadings of principal components analysis
513 axes using four inflorescence traits.


	PC 1	PC 2
Proportion of variance	54.11%	24.60%
Internode between bracteoles	−0.20	−0.96
Length of bracteole	0.61	−0.26
Width of bracteole	0.65	0.00
Style length	−0.41	0.08


514

515









Supplemental Table 1 Inflorescence/floral traits assessed on herbarium specimens. Inflorescence morphology types of Enclosing, Discoid-gland and Inconspicuous are shown as E, D and I, respectively. *N* indicates numbers of specimens observed (numbers of male and female specimens are shown separately). Presence/absence of disk-shaped glands on the middle part of the bracteole surfaces are indicated by “+” and “−”. Data not available are indicated by NA.

Species	Bracteole type	<i>N</i>		Disk-shaped glands	Length of bracteoles (mm)	Width of bracteoles (mm)	Internode between two bracteoles (mm)	Length of stigma (mm)
		♂	♀					
<i>M. albescens</i>	I	3	3	−	1.28	0.55	2.78	2.63
<i>M. alchorneoides</i>	I	3	0	−	1.14	1.12	2.76	NA
<i>M. aleuritoides</i>	D	3	3	+	1.19	0.39	2.00	6.43
<i>M. alnifolia</i>	D	2	2	+	1.43	0.37	3.58	4.37
<i>M. angustifolia</i>	D	3	2	+	4.19	2.18	5.23	0.34
<i>M. auriculata</i>	I	2	1	−	1.27	1.07	1.75	6.73
<i>M. barteri</i>	E	2	2	−	4.65	3.35	1.38	0.53
<i>M. bicolor</i>	E	4	0	−	2.33	2.17	0.99	NA
<i>M. bifoveata</i>	D	3	3	+	1.90	1.66	3.51	2.92
<i>M. brachytricha</i>	NA	0	0	NA	NA	NA	NA	NA
<i>M. clavata</i>	E	3	3	−	2.94	2.88	1.81	1.60
<i>M. conifera</i>	E	3	0	−	1.41	1.77	0.80	NA
<i>M. densiflora</i>	I	3	0	−	1.67	0.96	1.83	1.30
<i>M. denticulata</i>	D	5	4	+	1.01	1.03	NA	0.69
<i>M. diepenhorstii</i>	E	3	1	−	2.84	2.70	1.01	0.43
<i>M. domatiosa</i>	I	3	3	−	1.43	0.22	3.41	7.33

Supplemental Table 1 Continued.

Species	Bracteole type	N		Disk-shaped glands	Length of bracteoles (mm)	Width of bracteoles (mm)	Internode between two bracteoles (mm)	Length of stigma (mm)
		♂	♀					
<i>M. ducis</i>	I	2	0	—	2.43	1.27	1.88	NA
<i>M. echinocarpa</i>	NA	0	0	NA	NA	NA	NA	NA
<i>M. gabunica</i>	E	1	1	—	2.33	2.53	2.13	0.47
<i>M. gigantea</i>	E	3	0	—	3.40	3.43	1.91	NA
<i>M. grallata</i>	I	1	1	—	0.00	0.00	4.00	3.47
<i>M. grandifolia</i>	E	3	1	—	4.21	3.34	2.15	0.30
<i>M. heterophylla</i>	E	2	0	—	7.86	4.73	2.20	NA
<i>M. heynei</i>	E	4	0	—	5.69	4.39	1.97	NA
<i>M. hispida</i>	D	5	3	+	2.61	1.10	2.57	7.60
<i>M. hullettii</i>	E	3	0	—	3.61	2.69	2.07	NA
<i>M. hurifolia</i>	E	2	0	—	3.05	2.72	1.13	NA
<i>M. inamoena</i>	I	2	1	—	0.00	0.00	4.73	8.00
<i>M. indica</i>	D	5	2	+	3.19	1.33	7.30	2.03
<i>M. induta</i>	NA	0	2	—	NA	NA	NA	3.33
<i>M. involucrata</i>	D	2	2	+	0.97	0.60	2.20	2.77
<i>M. klaineana</i>	I	2	0	—	1.53	1.53	1.73	NA
<i>M. lamellata</i>	E	0	0	—	NA	NA	NA	NA
<i>M. lowii</i> var. <i>kostermansii</i>	I	3	2	—	1.39	0.72	1.56	6.57
<i>M. lowii</i> var. <i>lowii</i>	I	2	0	—	1.17	0.93	1.93	NA
<i>M. mauritiana</i>	I	2	0	—	2.15	2.20	3.55	NA

Supplemental Table 1 Continued.

Species	Bracteole type	N		Disk-shaped glands	Length of bracteoles (mm)	Width of bracteoles (mm)	Internode between two bracteoles (mm)	Length of stigma (mm)
		♂	♀					
<i>M. monandra</i>	E	3	0	—	4.71	3.42	1.22	NA
<i>M. novo-guineensis</i>	E	3	0	—	1.99	1.54	1.03	NA
<i>M. oblongifolia</i>	I	1	0	—	2.30	2.37	1.25	NA
<i>M. obovata</i>	D	2	1	+	1.85	0.80	1.75	3.60
<i>M. pachyphylla</i>	E	5	1	—	2.46	2.36	0.85	1.03
<i>M. pearsonii</i>	E	3	0	—	2.54	2.16	1.72	NA
<i>M. poggei</i>	E	2	0	—	2.30	2.37	1.25	NA
<i>M. puncticulata</i>	E	4	1	—	3.58	2.70	1.53	1.37
<i>M. praestans</i>	I	3	4	—	1.68	0.73	3.08	24.24
<i>M. quadriglandulosa</i>	E	3	3	—	2.72	2.42	1.23	3.40
<i>M. repando-dentata</i>	NA	0	3	NA	NA	NA	NA	38.11
<i>M. rhizinoides</i>	I	3	3	—	1.12	0.82	3.37	1.78
<i>M. saccifera</i>	E	1	0	—	2.97	3.17	1.13	NA
<i>M. schweinfurthii</i>	E	2	0	—	2.53	3.17	1.58	NA
<i>M. siamensis</i>	E	4	2	—	3.79	3.14	2.58	1.20
<i>M. sinensis</i>	D	3	3	+	3.17	1.09	3.46	0.68
<i>M. subdentata</i>	I	2	1	—	1.07	1.35	3.38	3.67
<i>M. tanarius</i>	E	3	3	—	6.00	4.97	2.98	3.83
<i>M. tessellata</i>	D	2	1	+	1.38	0.90	2.02	4.40
<i>M. trichocarpa</i>	E	4	3	—	4.28	2.63	1.51	0.88

Supplemental Table 1 Continued.

Species	Bracteole type	N		Disk-shaped glands	Length of bracteoles (mm)	Width of bracteoles (mm)	Internode between two bracteoles (mm)	Length of stigma (mm)
		♂	♀					
<i>M. triloba</i>	E	4	1	—	2.62	2.42	1.29	0.30
<i>M. umbrosa</i>	E	0	0	—	NA	NA	NA	NA
<i>M. winkleri</i>	E	4	3	—	2.83	2.70	2.08	0.26

Supplemental Figure 1 A female inflorescence of *Macaranga vedeliana*, not included in the analysis, classified into the Inconspicuous type.

